



Low-cost modified endoscopic vacuum therapy using a triple-lumen tube allows nutrition and drainage for treatment of an early post-bariatric surgery leak




► **Fig. 1** Underwater endoscopic visualization of the sleeve leak.

Leaks after laparoscopic sleeve gastrectomy (LSG) occur in up to 5% of patients, with mortality rates approaching 4% [1]. Among available therapies for leaks after bariatric surgery, endoscopic approaches have been shown to be safe and effective. Tissue sealants, cap-mounted clips, suturing, stents, a cardiac septal defect occluder, septotomy, endoscopic internal drainage with double pigtailed, and endoscopic vacuum therapy (EVT) are the forefront of the therapeutic endoscopic arsenal [2, 3]. However, an individualized approach is needed to determine the best endoscopic treatment strategy for a given patient. In this video, we describe the successful treatment of a LSG leak using low-cost modified EVT.

A 30-year-old woman with class II obesity developed fever, nausea, and abdominal pain 14 days after LSG. Computed tomographic scan revealed a leak associated with a small contained collection adjacent to the staple line of the proximal stomach (angle of His). Underwater esophagogastroduodenoscopy was performed without air insufflation to avoid disrupting the collection (► **Fig. 1**). The patient was treated with modified EVT using widely available triple-lumen tubing to allow for nutrition using the jejunal tube and EVT using the fenestrated portion of the tube (► **Video 1**). The procedure was well tolerated and the patient



► **Video 1** Low-cost modified endoscopic vacuum therapy with a triple-lumen tube.




► **Fig. 2** Fluoroscopic imaging confirmed the success of leak treatment with the low-cost modified endoscopic vacuum therapy (EVT) system and demonstrated a significant improvement in the sleeve stenosis.

remained on enteral nutrition as well as taking a liquid diet on the day following the procedure. External drainage was not required. After 15 days, imaging demonstrated resolution of the leak (► **Fig. 2**), and the modified EVT system was removed (► **Fig. 3**).

In summary, endoscopic drainage with this low-cost, modified EVT is a highly effective and safe technique for leaks [4,

5]. Furthermore, EVT obviates the need for external drainage and employs widely available materials. Ultimately, this low-cost modified EVT allows for nutrition and drainage with a single tube through the nares, enabling decreased procedure times, longer periods between EVT system exchanges, and fewer adverse events.







► **Fig. 3** Complete healing of the leak after one session of the modified EVT treatment.

Endoscopy_UCTN_Code_TTT_1AO_2AI

Competing interests

The authors declare that they have no conflict of interest.

The authors

Diogo Turiani Hourneaux de Moura¹ , **Bruno Salomão Hirsch**¹ , **Mateus Bond Boghossian**¹ , **Flaubert Sena de Medeiros**², **Thomas R. McCarty**³, **Christopher C. Thompson**³, **Eduardo Guimarães Hourneaux de Moura**¹ 

- 1 Gastrointestinal Endoscopy Unit, Gastroenterology Department, Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil
- 2 Surgery Division, Federal University of Rio Grande do Norte, Natal, Brazil
- 3 Gastroenterology, Hepatology and Endoscopy Division – Harvard Medical School, Brigham and Women’s Hospital, Boston, Massachusetts, USA

Corresponding author

Bruno Salomão Hirsch, MD
Av. Dr Enéas de Carvalho Aguiar, 225,
6º andar, bloco 3, Cerqueira César,
05403-010 – São Paulo, SP, Brazil
brunosalomah@hotmail.com

References

- [1] Hamid HKS, Emile SH, Saber AA et al. Customized bariatric stents for sleeve gastrectomy leak: are they superior to conventional esophageal stents? A systematic review and proportion meta-analysis *Surg Endosc* 2021; 35: 1025–1038
- [2] de Moura DTH, Sachdev AH, Thompson CC. Endoscopic full-thickness defects and closure techniques. *Curr Treat Options Gastroenterol* 2018; 16: 386–405
- [3] do Monte ES Jr, de Moura DTH, Ribeiro IB et al. Endoscopic vacuum therapy versus endoscopic stenting for upper gastrointestinal transmural defects: systematic review and meta-analysis. *Dig Endosc* 2020. doi:10.1111/den.13813
- [4] Loske G, Schorsch T, Rucktaeschel F et al. Open-pore film drainage (OFD): a new multipurpose tool for endoscopic negative pressure therapy (ENPT). *Endosc Int Open* 2018; 6: E865–E871
- [5] de Moura DTH, de Moura BFBH, Manfredi MA et al. Role of endoscopic vacuum therapy in the management of gastrointestinal transmural defects. *World J Gastrointest Endosc* 2019; 11: 329–344

Bibliography

Endoscopy
DOI 10.1055/a-1540-5870
ISSN 0013-726X
published online 2021
© 2021. Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>